

# Molecular enhancement of porcine cardiac chronotropy

J M Edelberg, D T Huang, M E Josephson, R D Rosenberg

## Abstract

**Objective**—To test the potential of gene transfer approaches to enhance cardiac chronotropy in a porcine system as a model of the human heart.

**Methods**—Plasmids encoding either the human  $\beta_2$  adrenergic receptor or control constructs were injected into the right atria of native Yorkshire pig hearts. Percutaneous electrophysiological recording catheters equipped with 33 gauge circular injection needles were positioned in the mid-lateral right atrium. At the site of the earliest atrial potential the circular injection needles were rotated into the myocardium and the  $\beta_2$  adrenergic receptor (n = 6) or control plasmid constructs (n = 5) were injected.

**Results**—Injection of the  $\beta_2$  adrenergic receptor construct significantly enhanced chronotropy compared with control injections. The average (SD) heart rate of the pigs was 108 (16) beats/min before injection. Two days after injection with control plasmids the heart rate was 127 (25) beats/min (NS compared with preinjection rates). After injection with plasmid encoding the  $\beta_2$  adrenergic receptor the heart rate increased by 50% to 163 (33) beats/min ( $p < 0.05$  compared with preinjection and postinjection control rates).

**Conclusions**—The present studies showed in a large animal model that local targeting of gene expression may be a feasible modality to regulate cardiac pacemaking activity. In addition, these investigations provide an experimental basis for developing future clinical gene transfer approaches to upregulate heart rate and modulate cardiac conduction.

(Heart 2001;86:559–562)

Keywords: sinus node; adrenergic receptor; chronotropic agents; conduction system; gene therapy

Cardiac chronotropic incompetence is associated with increased morbidity and mortality.<sup>1–3</sup> Permanent chronotropic disorders caused by sinus node diseases include inappropriate sinus bradycardia, sinus arrest, and sinus exit block.<sup>4–5</sup> These diseases are not confined to the aged, being occasionally diagnosed in young adults,<sup>6–7</sup> and may have a familial component.<sup>8–10</sup> The pathology of the disease is diverse. Anatomical studies have shown the presence of fibrosis and sclerodermal infiltrates in the sinus node associated with advanced age.<sup>11–12</sup> Ischaemic and infectious causes have also been linked to chronotropic incompetence.<sup>13–15</sup> The treatment of these disorders is limited, with only a minority of the transient disorders amenable to medical treatment.<sup>16</sup> The majority of the causes of chronotropic incompetence require the implantation of an electronic pacemaker, either temporarily or permanently.<sup>17</sup> Future treatments for chronotropic incompetence may be based on therapeutics that can specifically enhance the pacemaker potential of the endogenous cardiac tissue and potentially diminish the necessity for exogenous electronic devices.

Recent advances in biomedical research suggest that gene therapy may be used in the treatment of cardiovascular disease.<sup>18–19</sup> Cardiac pacemaking offers a unique target in gene therapy as modulation of the spontaneous depolarisations in a selected set of cells may enhance heart rate. Recently we reported the development of a murine gene transfer model to test the effects of candidate genes on heart rate.<sup>20</sup> In this model constructs encoding the

human  $\beta_2$  adrenergic receptor were used under in vitro, ex vivo, and in vivo conditions to enhance selectively murine cardiac chronotropy, suggesting a potential role for this approach in future treatments of cardiac pacemaking disorders.

The translation into clinical treatment requires that this strategy be further tested and optimised in large animal models. The Yorkshire pig model was used in the present study based on its anatomical and physiological similarity to the human cardiovascular system.<sup>21</sup> These studies offered the opportunity to develop a transvenous catheter delivery gene transfer approach to enhance cardiac chronotropy, which may be adapted for future human investigations.

## Methods

### PLASMID CONSTRUCTS

A 2.25 kb *SalI*-*Bam*HI fragment, the human  $\beta_2$  adrenergic receptor complementary DNA (cDNA), was ligated into a *SalI*-*Bam*HI site 3' to the  $\beta$  actin promoter in a pBR322 vector to generate pBR322- $\beta$  actin promoter- $\beta_2$  adrenergic receptor. In similar fashion, the bacterial  $\beta$  galactosidase gene was ligated to the  $\beta$  actin promoter in a pBR322 vector and served as a control vector. The plasmid construct encoding the humanised green fluorescent protein<sup>22</sup> with a cytomegalovirus promoter element was purchased from Clontech (Palo Alto, California, USA) and served as a co-injection vector. The recombinant DNA injection solutions (200  $\mu$ l; 100  $\mu$ g DNA/ml; 5:1 M/M  $\beta_2$  adrenergic receptor or  $\beta$  galactosidase/green fluorescent protein)

Department of  
Medicine, Beth Israel  
Deaconess Medical  
Center, Boston, MA  
02215, USA

J M Edelberg  
D T Huang  
M E Josephson

Department of  
Biology, Massachusetts  
Institute of  
Technology,  
Cambridge, MA 02139,  
USA  
R D Rosenberg

Correspondence to:  
Dr Edelberg  
jme2002@  
mail.med.cornell.edu

Accepted 11 July 2001

were prepared with in an injection vehicle of phosphate buffered saline (PBS) with 20% sucrose and 2% Evans blue.

#### INJECTION CATHETER

The electrophysiological recording catheter was custom designed and manufactured by Medtronic, Inc (Minneapolis, Minnesota, USA). The polyurethane coated catheter was 7 French gauge and was supported with an 8 French sheath. The distal end of the catheter was terminated with a three and a half turn 33 gauge corkscrew shaped needle allowing it to impale tissues securely to record local intracardiac electrograms. The proximal end of the catheter was terminated with a luer lock injection port allowing it to accept standard sized syringes.

#### NATIVE HEART DNA INJECTION

Female Yorkshire pigs weighing 15–20 kg were initially anaesthetised with intramuscular ketamine (10 g/kg) and intubated. The animals were then given 2% isoflurane and ventilated with a large animal ventilator (Hallowell model 2000, Pittsfield, MA, USA). Heart rate, blood pressure, and arterial oxygen saturation were monitored for the duration of the procedure. By sterile technique, the right femoral vein was exposed and cannulated, and an 8 French sheath was inserted. Under fluoroscopic guidance the electrophysiological injection catheter was introduced and advanced to the right atrium. Simultaneous six lead surface and intracardiac electrocardiograms were recorded with a multichannel recorder (EVR PPG Biomedical, Pittsburgh, PA, USA). The A-P interval (ms), cycle length (ms), PR interval (ms), and P wave axis ( $^{\circ}$ ) were measured. At the site of the earliest A wave the injection needle was rotated  $270^{\circ}$  into the atrial myocardium. The recombinant DNA construct (200  $\mu$ l) controls ( $n = 5$ ) or  $\beta_2$  adrenergic receptor samples ( $n = 6$ ) were then injected into the atrial myocardium in a randomised double blinded protocol. The catheter was then disengaged and removed from the animal. The animals were observed for an additional 10 min and monitored for complications. The vascular sheath was then removed, the vein was sutured, and the incision site was closed. Anaesthesia was then discontinued. After they regained spontaneous respirations the animals were placed in individual pens. The animals were monitored hourly for the next 3 h and then daily until termination of the experiments, 48–96 h after injection.

#### SERIAL SURFACE ECGS AND ANALYSIS

Serial surface ECGs were recorded daily for all animals for the duration of the study. The pigs were anaesthetised with ketamine as above. Simultaneous six lead surface ECGs were recorded. The cycle length, PR interval, and P wave axis were measured (mean (SD)). Significance was determined by a Student's *t* test analysis between average preinjection and postinjection heart rates with the control and  $\beta_2$  adrenergic receptor plasmid injections, and between the percentage change in paired

preinjection and postinjection heart rates with the control and  $\beta_2$  adrenergic receptor plasmid injections.


#### $\beta_2$ ADRENERGIC RECEPTOR IMMUNOSTAINING

At the termination of the experiments, the animals were sacrificed and the hearts explanted. The injection sites were identified by Evans blue and harvested unfixed for cryosectioning and immunostaining. The samples were cut in to 10  $\mu$ m sections and fixed with cold methanol for 10 minutes. The sections were then washed with PBS and blocked with 10% normal serum in PBS for 20 minutes. Samples were then incubated with a rabbit antihuman  $\beta_2$  adrenergic receptor polyclonal antibody (Santa Cruz Biotechnologies, Santa Cruz, California, USA; 1.0  $\mu$ g/ml) for one hour in a humid chamber at  $25^{\circ}\text{C}$ . The samples were washed with PBS three times and then incubated with a secondary donkey antirabbit Cy3 polyclonal antibody (Jackson ImmunoResearch, West Grove, Pennsylvania, USA) at a 1:1000 dilution in PBS with 1% bovine serum albumin for one hour in a humid chamber at room temperature. The samples were washed with PBS three times and mounted with 90% glycerol in PBS. Green fluorescent protein expression was identified by using epifluorescence filters (excitation 405 nm/emission 490 nm). Immunostaining for the human  $\beta_2$  adrenergic receptor was identified by using epifluorescence filters (excitation 488 nm/emission 540 nm).

#### Results

The animals were anaesthetised and intubated, and venous access was obtained as described above. The injection catheter was advanced to the right lateral atrium under fluoroscopic guidance (fig 1A). Simultaneous surface and intracardiac ECGs were recorded. The catheter was positioned at the site of the earliest atrial activity, as fig 1B shows. The atrial potential at the injection sight was similar in the pigs injected with the control (14 (10) ms) and the  $\beta_2$  adrenergic receptor encoding constructs (12 (10) ms). In addition, both the average PR interval and the P wave axis on the surface ECG were similar in the groups before injection (table 1). All the animals tolerated the procedure well.

Serial surface ECGs recorded from the pigs after construct injection showed that the average PR interval and P wave axis were similar to the measurements before injection (table 1). The heart rate increased in the hearts injected with the  $\beta_2$  adrenergic receptor plasmid compared to hearts with the control injections (fig 2A). The average heart rate of the pigs was 108 (16) beats/min before injection. Two days after injection with control plasmids the heart rate was 127 (25) beats/min ( $p > 0.3$  compared with preinjection). After injection with plasmid encoding the  $\beta_2$  adrenergic receptor the heart rate increased to 163 (33) beats/min ( $p < 0.05$  compared with preinjection), an approximately 50% increase in heart rate ( $p < 0.01$  compared with the change in postinjection control heart rate) (fig 2B). All



**Figure 1** Cine of electrophysiological recording and injection catheter (arrow head) during injection of complementary DNA in to the porcine lateral right atrium (A). Two examples are shown of surface and intracardiac ECGs recorded before injection into the right atrium (B).

**Table 1** Electrocardiographic measurements

	Baseline		48 h postinjection	
	Control	$\beta_2$ AR	Control	$\beta_2$ AR
P wave axis ( $^{\circ}$ )	65 (7)	56 (20)	60 (20)	62 (15)
PR interval (ms)	85 (10)	92 (9)	86 (8)	86 (9)


$\beta_2$ AR,  $\beta_2$  adrenergic receptor.

animals survived until the termination of the experiments.

Injection of the cDNA constructs led to the expression of the encoded genes. Sections of the right atrial tissue at the site of injection showed the presence of green fluorescent protein (fig 2C,D). Immunostaining of the injection site sections showed colocalisation of the human  $\beta_2$  adrenergic receptor in the hearts co-injected with the  $\beta_2$  adrenergic receptor encoding constructs (fig 2D) but not in those injected with control constructs (fig 2C).

## Discussion

The present experiments were directed at developing in vivo gene transfer techniques to identify and study genes that can be used to upregulate heart rate selectively and alter cardiac rhythm in the intact heart in a large animal model. Our goal is eventually to develop a molecular or cellular based approach to the treatment of cardiac chronotropic and conduction disorders. We have focused our initial efforts on a direct porcine cardiac gene transfer



**Figure 2** (A) Representative surface ECGs recorded 48 hours after injection of either control constructs or constructs encoding  $\beta_2$  adrenergic receptor. (B) The average percentage change in heart rate (HR) after injection of control construct or  $\beta_2$  adrenergic receptor ( $\beta_2$ AR) encoding plasmids (\* $p < 0.01$  versus control). (C) Dual fluorescence micrographs of sections of right atrial tissue co-injected with plasmids encoding green fluorescent protein and control constructs (C) and with the human  $\beta_2$  adrenergic receptor (D). Green fluorescent protein was visualised directly (green), and the human  $\beta_2$  adrenergic receptor was detected by immunostaining (red). Bar 32  $\mu$ m.

system to translate our recently developed murine cDNA chronotropic test system<sup>20</sup> into a clinically applicable endovascular therapeutic approach.

The transvenous delivery of the cDNA constructs was electrophysiologically guided. Constructs were injected into the area of the sinus node in an attempt to enhance the chronotropy of the intact cardiac conduction system. The mappings of the perisinus node injection sites were in agreement with previous characterisations of the porcine sinus node.<sup>23</sup> Delivery of the cDNA encoding the human  $\beta_2$  adrenergic receptor resulted in a significant increase in the heart rate compared with control injections. The non-significant change in the control injection heart rate may have been caused by the level of anaesthesia or the intubation of the animals at baseline, as has previously been shown in porcine anaesthesia studies.<sup>24,25</sup>

The  $\beta_2$  adrenergic receptor injection enhancement of chronotropy was not associated with altered characteristics of atrial conduction, with no significant changes in either the PR interval or the P wave axis. These data suggest that the increased heart rate is mediated by

an increased rate of sinus node depolarisations. Alternatively, the enhanced heart rate may arise from a perisinus nodal atrial focus that conducts normally through the atrium. Future approaches will target atrial sites removed from the sinus node, including the left atrium, to test the potential of gene transfer to establish independent pacemaking foci.

The transience of the increase in porcine heart was expected given the use of plasmid cDNA constructs. Previous research has shown that injection of plasmid into the left ventricular myocardium may result in prolonged construct expression (> 1–2 months)<sup>26, 27</sup> compared with the 2–3 days observed in the murine<sup>20</sup> and the present porcine right atrial injections. The limited duration seen in the present study is likely multifactorial. Previous studies have shown significant differences in the expression of plasmid constructs injected into different muscle groups.<sup>28, 29</sup> These regional variations may have an anatomical basis as a result of technical injection limitations as well as variable DNA diffusion rates in various tissues.<sup>28</sup> Additional factors, including atrial T tubule density,<sup>30, 31</sup> may also account for the relatively short period of plasmid mediated enhancement of cardiac chronotropy, and different vector strategies will be required to sustain the increased heart rates. To this end, advances in adenoviral and adenoviral associated viral technologies offer efficient approaches for more extended expression of constructs in cardiac tissues.<sup>32–34</sup> Moreover, cardiac myocyte transplantation approaches<sup>35, 36</sup> may allow for the engraftment of a more permanent genetically engineered tissue to modulate the cardiac conduction system.

In summary, these studies showed that the basal rate of the heart can be enhanced by local intravascular delivery of exogenous genes. These investigations may provide a foundation for developing novel therapeutics for cardiac chronotropic disorders, such as sick sinus syndrome.

cDNA encoding the human  $\beta_1$  adrenergic receptor was the kind gift of Dr Robert J Lefkowitz (Duke University Medical Center, Durham, NC, USA). This work was done during the tenure of a research fellowship from the American Heart Association, Massachusetts Affiliate, Inc. (to JME). This work was supported by National Institutes of Health Grants HL41484 and HL59316 (RDR) and American Heart Association Fellowship Award #13-419-967 (JME).

- Brignole M, Menozzi C, Lolli G, et al. Pacing for carotid sinus syndrome and sick sinus syndrome. *Pacing Clin Electrophysiol* 1990;13:2071–5.
- Ogawa H, Inoue T, Miwa S, et al. Heart rate responses to autonomic drugs in sick sinus syndrome: correlation with syncope and electrophysiologic data. *Jpn Circ J* 1991;55:15–23.
- Rodriguez RD, Schocken DD. Update on sick sinus syndrome, a cardiac disorder of aging. *Geriatrics* 1990;45:26–30, 33–6.
- Asseman P, Berzin B, Desry D, et al. Postextrasystolic sinoatrial exit block in human sick sinus syndrome: demonstration by direct recording of sinus node electrograms. *Am Heart J* 1991;122:1633–43.
- Wu DL, Yeh SJ, Lin FC, et al. Sinus automaticity and sinoatrial conduction in severe symptomatic sick sinus syndrome. *J Am Coll Cardiol* 1992;19:355–64.
- Kay R, Estioko M, Wiener I. Primary sick sinus syndrome as an indication for chronic pacemaker therapy in young adults: incidence, clinical features, and long-term evaluation. *Am Heart J* 1982;103:338–42.
- Yabek SM, Dillon T, Berman W, et al. Symptomatic sinus node dysfunction in children without structural heart disease. *Pediatrics* 1982;69:590–3.
- Olofsson BO, Eriksson P, Eriksson A. The sick sinus syndrome in familial amyloidosis with polyneuropathy. *Int J Cardiol* 1983;4:71–3.
- Onat A. Familial sinus node disease and degenerative myopia: a new hereditary syndrome? *Hum Genet* 1986;72:182–4.
- Mehta AV, Chidambaram B, Garrett A. Familial symptomatic sinus bradycardia: autosomal dominant inheritance. *Pediatr Cardiol* 1995;16:231–4.
- Shiraishi I, Takamatsu T, Minamikawa T, et al. Quantitative histological analysis of the human sinoatrial node during growth and aging. *Circulation* 1992;85:2176–84.
- Bharati S, Lev M. The pathologic changes in the conduction system beyond the age of ninety. *Am Heart J* 1992;124:486–96.
- Shaw DB, Linker NJ, Heaver PA, et al. Chronic sinoatrial disorder (sick sinus syndrome): a possible result of cardiac ischaemia. *Br Heart J* 1987;58:598–607.
- Junqueira Junior LF, Beraldo PS, Chapadeiro E, et al. Cardiac autonomic dysfunction and neuroganglionitis in a rat model of chronic Chagas' disease. *Cardiovasc Res* 1992;26:324–9.
- Molina HA, Milei J, Rimoldi MT, et al. Histopathology of the heart conducting system in experimental Chagas disease in mice. *Trans R Soc Trop Med Hyg* 1988;82:241–6.
- Alboni P, Menozzi C, Brignole M, et al. Effects of permanent pacemaker and oral theophylline in sick sinus syndrome the THEOPACE study: a randomized controlled trial. *Circulation* 1997;96:260–6.
- Alagona P Jr. Advances in pacing for the patient with sick sinus syndrome. *Curr Opin Cardiol* 1997;12:3–11.
- Finkel T, Epstein SE. Gene therapy for vascular disease. *FASEB J* 1995;9:843–51.
- Libby P. Gene therapy of restenosis: promise and perils. *Circ Res* 1998;82:404–6.
- Edelberg JM, Aird WC, Rosenberg RD. Enhancement of murine cardiac chronotropy by the molecular transfer of the human  $\beta_2$  adrenergic receptor cDNA. *J Clin Invest* 1998;101:337–43.
- Hughes HC. Swine in cardiovascular research. *Lab Anim Sci* 1986;36:348–50.
- Haas J, Park EC, Seed B. Codon usage limitation in the expression of HIV-1 envelope glycoprotein. *Curr Biol* 1996;6:315–24.
- Gepstein L, Hayam G, Ben-Haim SA. A novel method for nonfluoroscopic catheter-based electroanatomical mapping of the heart: in vitro and in vivo accuracy results. *Circulation* 1997;95:1611–22.
- Sundeman H, Haney M, Broome M, et al. The effects of desflurane on cardiac function as measured by conductance volumetry in swine. *Anesth Analg* 1998;87:522–8.
- Heerdts PM, Gandhi CD, Dickstein ML. Disparity of isoflurane effects on left and right ventricular afterload and hydraulic power generation in swine. *Anesth Analg* 1998;87:511–21.
- Lin H, Parmacek MS, Morle G, et al. Expression of recombinant genes in myocardium in vivo after direct injection of DNA. *Circulation* 1990;82:2217–21.
- Buttrick PM, Kass A, Kitsis RN, et al. Behavior of genes directly injected into the rat heart in vivo. *Circ Res* 1992;70:193–8.
- Wolff JA, Williams P, Acsadi G, et al. Conditions affecting direct gene transfer into rodent muscle in vivo. *Biotechniques* 1991;11:474–85.
- von Harsdorf R, Schott RJ, Shen YT, et al. Gene injection into canine myocardium as a useful model for studying gene expression in the heart of large mammals. *Circ Res* 1993;72:688–95.
- Wolff JA, Dowty ME, Jiao S, et al. Expression of naked plasmids by cultured myotubes and entry of plasmids into T tubules and caveolae of mammalian skeletal muscle. *J Cell Sci* 1992;103:1249–59.
- Tidball J, Cederdahl J, Bers D. Quantitative analysis of regional variability in the distribution of transverse tubules in rabbit myocardium. *Cell Tissue Res* 1991;264:293–8.
- Kass-Eisler A, Falck-Pedersen E, Alvira M, et al. Quantitative determination of adenovirus-mediated gene delivery to rat cardiac myocytes in vitro and in vivo. *Proc Natl Acad Sci U S A* 1993;90:11498–502.
- Lee J, Laks H, Drinkwater DC, et al. Cardiac gene transfer by intracoronary infusion of adenovirus vector-mediated reporter gene in the transplanted mouse heart. *J Thorac Cardiovasc Surg* 1996;111:246–52.
- Johns DC, Nuss HB, Chiamvimonvat N, et al. Adenovirus-mediated expression of a voltage-gated potassium channel in vitro (rat cardiac myocytes) and in vivo (rat liver): a novel strategy for modifying excitability. *J Clin Invest* 1995;96:1152–8.
- Koh GY, Soonpaa MH, Klug MG, et al. Stable fetal cardiomyocyte grafts in the hearts of dystrophic mice and dogs. *J Clin Invest* 1995;96:2034–42.
- Aoki M, Morishita R, Higaki J, et al. Survival of grafts of genetically modified cardiac myocytes transfected with FITC-labeled oligodeoxynucleotides and the beta-galactosidase gene in the noninfarcted area, but not the myocardial infarcted area. *Gene Ther* 1997;4:120–7.